

**REMARKS**

Attached hereto is a marked-up version of the changes made to the specification, sequence listing and claims by the current amendment. The attached marked-up version is captioned "**Version with markings to show changes made.**"

Should the Examiner have any questions, the Examiner is encouraged to telephone the undersigned.

Respectfully submitted,

Date: Oct. 29, 2001

By: David J. Weitz  
David J. Weitz  
Registration No. 38,362

WILSON SONSINI GOODRICH & ROSATI  
650 Page Mill Road  
Palo Alto, CA 94304-1505  
(650) 493-9300  
Customer No. 021971

RECEIVED

VERSION WITH MARKINGS TO SHOW CHANGES MADE

In the Specification:

The paragraph beginning at line 8, page 1 has been amended as follows:

[This application is a continuation-in-part of "ENERGY TRANSFER DYES WITH ENHANCED FLUORESCENCE," Application Serial No.: 08/642,330; Filed: May 3, 1996 and U.S. Application Serial No.: 08/672,196; filed June 27, 1996; entitled: "4,7-DICHLORORHODAMINE DYES" which are incorporated herein by reference.]

This application is a continuation of U.S. Application Serial No. 09/272,097, filed on March 18, 1999, which is a continuation of U.S. Application Serial No. 09/046,203, filed on March 23, 1998, now Patent No. 5,945,526, which is a continuation of U.S. Application No. 08/726,462, filed on October 4, 1996, now Patent No. 5,800,996, which is a continuation-in-part of U.S. Application No. 08/672,196, filed on June 27, 1996, now U.S. Patent No. 5,847,162 which is a continuation-in-part of U.S. Application No. 08/642,330, filed on May 3, 1996, now Patent No. 5,863,727, each of which it is incorporated herein by reference in its entirety.

The paragraph beginning at line 20, on page 10, has been amended as follows:

In another embodiment, the energy transfer fluorescent dyes have donor and acceptor dyes with the general structure where  $Y_1$  and  $Y_2$  taken separately are either hydroxyl, oxygen, iminium or amine, the iminium and amine preferably being a tertiary iminium or amine and  $[R_{11}-R_{17}] R_{11}-R_{16}$  are any substituents which are compatible with the energy transfer dyes of the present invention.

Page 52 has been amended as follows:

In compound [3a] 3A-A, one of  $R_1$  and  $R_2$  is ethyl, the other being hydrogen,  $R_3$  and  $R_4$  taken separately are hydrogen,  $[R_5]$   $R_6$  is methyl,  $[R_6-R_{10}]$   $R_5$  and  $R_7-R_{10}$  taken separately

are hydrogen,  $X_1$  is carboxylate, and one of  $X_3$  and  $X_4$  is a linking group, the other being hydrogen.

In compound [3b] 3A-B, one of  $R_1$  and  $R_2$  is ethyl, the other being hydrogen,  $R_3$  and  $R_4$  taken separately are methyl,  $R_5$  is methyl,  $R_6$ - $R_{10}$  taken separately are hydrogen,  $X_1$  is carboxylate, and, one of  $X_3$  and  $X_4$  is a linking group, the other being hydrogen.

In compound [3c] 3A-C,  $R_1$  and  $R_2$  taken separately are methyl,  $R_3$  and  $[R_7] R_9$  taken together form a six membered ring,  $R_4$  and  $R_8$  taken together form a six membered ring,  $R_5$ ,  $R_6$ ,  $[R_9] R_2$ , and  $R_{10}$  taken separately are hydrogen,  $X_1$  is carboxylate, and, one of  $X_3$  and  $X_4$  is a linking group, the other being hydrogen.

In compound [3d] 3B-D,  $R_1$  and  $R_2$  taken separately are hydrogen,  $R_3$  and  $[R_7] R_9$  taken together form a six membered ring,  $R_4$  and  $R_8$  taken together form a six membered ring,  $R_5$ ,  $R_6$ ,  $[R_9] R_2$ , and  $R_{10}$  taken separately are hydrogen,  $X_1$  is carboxylate, and one of  $X_3$  and  $X_4$  is a linking group, the other being hydrogen.

In compound [3e] 3B-E, one of  $R_1$  and  $R_2$  is ethyl, the other being hydrogen,  $R_3$  and  $[R_7] R_9$  taken together form a six membered ring,  $R_4$  and  $R_8$  taken together form a six membered ring,  $R_5$  is methyl,  $R_6$ ,  $[R_9] R_2$  and  $R_{10}$  taken separately are hydrogen,  $X_1$  is carboxylate, and, one of  $X_3$  and  $X_4$  is a linking group, the other being hydrogen.

In compound [3f] 3B-F,  $R_1$  and  $R_2$  taken separately are hydrogen,  $R_3$  and  $R_4$  taken separately are methyl,  $R_5$ - $R_{10}$  taken separately are hydrogen,  $X_1$  is carboxylate, and, one of  $X_3$  and  $X_4$  is linking group, the other being hydrogen.

The paragraph beginning at line 5, on page 53 has been amended as follows:

Figure 4A shows a generalized synthesis wherein the substituent  $X_1$  can be other than carboxylate. In the figure,  $X'$  indicates moieties which are precursors to  $X_1$ . In the method illustrated in Figure 4A, two equivalents of a 3-aminophenol derivative [4a/4b] 4A-A/4A-B, such as 3-dimethylaminophenol, is reacted with one equivalent of a dichlorobenzene derivative [4c] 4A-C, e.g., 4-carboxy-3,6,dichloro-2-sulfobenzoic acid cyclic anhydride, i.e., where the  $X_1'$  moieties of 4c taken together are,

The paragraphs beginning at line 14, on page 53, have been amended as follows:

The reactants are then heated for 12 h in a strong acid, e.g., polyphosphoric acid or sulfuric acid, at 180°C. The crude dye [4d] 4A-D is precipitated by addition to water and isolated by centrifugation. To form a symmetrical product, the substituents of reactants [4a] 4A-A and 4b are the same, while to form an asymmetrical product, the substituents are different.

Figure 4B shows a generalized synthesis wherein the substituent X<sub>1</sub> is carboxylate. In the method of Figure 4B, two equivalents of a 3-aminophenol derivative [4a/4b] 4A-A/4A-B, such as 3-dimethylaminophenol, is reacted with one equivalent of a phthalic anhydride derivative [4e] 4B-E, e.g. 3,6-dichlorotrimellitic acid anhydride. The reactants are then heated for 12 h in a strong acid, e.g., polyphosphoric acid or sulfuric acid, at 180°C. The crude dye [4d] 4A-D is precipitated by addition to water and isolated by centrifugation. To form a symmetrical product, the substituents of reactants [4a] 4A-A and [4b] 4A-B are the same, while to form an asymmetrical product, the substituents are different.

The paragraph beginning at line 23, page 93, has been amended as follows:

Dye primer sequencing was performed on the [M13 (SEQ. ID. NO.: 2)] pGEM (SEQ. ID. NO.: 3) using a set of four dyes attached to the M13-21 primer (SEQ. ID. [NO. 3] NO.: 2) as described in Example 5. Figure 13 is a four color plot of the dye labeled oligonucleotides produced from the sequencing. The peak for cytosine corresponds to the fluorescence of 5-CFB-DR110-2. The peak for adenosine corresponds to the fluorescence of 6-CFB-DR6g-2. The peak for guanosine corresponds to the fluorescence of 5-CFB-DTMR-2. The peak for thymidine corresponds to the fluorescence of 5-CFB-DROX-2.